

# CAUSES OF INFANT DEATHS

*2012 - 13*

The HMIS data is taken from National Web Portal

**NHSRC**

**9/23/2013**



## India - Causes of Infant Deaths - HMIS - 2012 - 13

Estimated Population - 2012	1,230,384,560	
CBR - SRS - 2012	21.8	
Estimated Live Births	26,822,383	
Reported Infant Deaths	160,003	
Infant deaths per 1000 Estimated live Births	6	
Early Neonatal deaths per 1000 Estimated live Births	4	
Late Neonatal deaths per 1000 Estimated live Births	1	
	Total	%ge against Total Infant Deaths reported
Number of cases of Infant deaths within 24 hrs of birth	23,739	15%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	6,214	4%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	2,909	2%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	15,829	10%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	2,573	2%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	23,953	15%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	6,898	4%

India - Causes of Infant Deaths - HMIS - 2012 - 13		
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	27,345	17%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	12,861	8%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	6,969	4%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	1,610	1%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	3,132	2%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	242	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	25,729	16%

Andhra Pradesh - Causes of Infant Deaths - HMIS - 2012 - 13	
Estimated Population - 2012	85,615,407
CBR - SRS - 2012	17.5
Estimated Live Births	1,498,270
Reported Infant Deaths	10,477
Infant deaths per 1000 Estimated live Births	7

## Andhra Pradesh - Causes of Infant Deaths - HMIS - 2012 - 13

Early Neonatal deaths per 1000 Estimated live Births	3.38	
Late Neonatal deaths per 1000 Estimated live Births	1.77	
	<b>Total</b>	<b>%ge against Total Infant Deaths reported</b>
Number of cases of Infant deaths within 24 hrs of birth	343	3%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	333	3%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	148	1%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	688	7%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	221	2%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	1,610	15%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	676	6%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	2,093	20%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	1,602	15%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	258	2%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	27	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever	129	1%

### Andhra Pradesh - Causes of Infant Deaths - HMIS - 2012 - 13

related		
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	31	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	2,318	22%

### Arunachal Pradesh - Causes of Infant Deaths - HMIS - 2012 - 13

Estimated Population - 2012	1,415,738	
CBR - SRS - 2012	19.8	
Estimated Live Births	28,032	
Reported Infant Deaths	50	
Infant deaths per 1000 Estimated live Births	2	
Early Neonatal deaths per 1000 Estimated live Births	0.96	
Late Neonatal deaths per 1000 Estimated live Births	0.04	
	Total	%ge against Total Infant Deaths reported

### Arunachal Pradesh - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant deaths within 24 hrs of birth	3	6%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	-	0%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	-	0%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	-	0%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	-	0%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	1	2%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	-	0%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	23	46%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	1	2%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	6	12%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	6	12%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Fever related	1	2%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Measles	-	0%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	9	18%

Assam - Causes of Infant Deaths - HMIS - 2012 - 13		
Estimated Population - 2012	31,662,740	
CBR - SRS - 2012	22.8	
Estimated Live Births	721,910	
Reported Infant Deaths	5,346	
Infant deaths per 1000 Estimated live Births	7	
Early Neonatal deaths per 1000 Estimated live Births	4.28	
Late Neonatal deaths per 1000 Estimated live Births	0.70	
	Total	%ge against Total Infant Deaths reported
Number of cases of Infant deaths within 24 hrs of birth	330	6%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	196	4%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	10	0%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	692	13%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	21	0%

### Assam - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	692	13%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	47	1%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	1,183	22%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	425	8%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	313	6%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	61	1%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	215	4%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	20	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	1,141	21%



Bihar - Causes of Infant Deaths - HMIS - 2012 - 13		
Estimated Population - 2012	106,168,110	
CBR - SRS - 2012	27.7	
Estimated Live Births	2,940,857	
Reported Infant Deaths	3,902	
Infant deaths per 1000 Estimated live Births	1	
Early Neonatal deaths per 1000 Estimated live Births	0.85	
Late Neonatal deaths per 1000 Estimated live Births	0.14	
	Total	%ge against Total Infant Deaths reported
Number of cases of Infant deaths within 24 hrs of birth	1,164	30%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	230	6%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	77	2%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	367	9%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	45	1%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	271	7%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	46	1%

Bihar - Causes of Infant Deaths - HMIS - 2012 - 13		
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	468	12%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	257	7%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	171	4%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	170	4%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	278	7%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	108	3%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	250	6%

Chhattisgarh - Causes of Infant Deaths - HMIS - 2012 - 13	
Estimated Population - 2012	26,070,464
CBR - SRS - 2012	24.9
Estimated Live Births	649,155

## Chhattisgarh - Causes of Infant Deaths - HMIS - 2012 - 13

<b>Reported Infant Deaths</b>	3,977	
<b>Infant deaths per 1000 Estimated live Births</b>	6	
<b>Early Neonatal deaths per 1000 Estimated live Births</b>	3.67	
<b>Late Neonatal deaths per 1000 Estimated live Births</b>	0.40	
	<b>Total</b>	<b>%ge against Total Infant Deaths reported</b>
<b>Number of cases of Infant deaths within 24 hrs of birth</b>	366	9%
<b>Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis</b>	117	3%
<b>Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis</b>	19	0%
<b>Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia</b>	248	6%
<b>Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia</b>	10	0%
<b>Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW</b>	920	23%
<b>Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW</b>	121	3%
<b>Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW</b>	732	18%
<b>Number of cases of Infant deathsone week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW</b>	107	3%
<b>Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia</b>	411	10%

### Chhattisgarh - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	35	1%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	96	2%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	1	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	794	20%

### Goa - Causes of Infant Deaths - HMIS - 2012 - 13

Estimated Population - 2012	1,486,877	
CBR - SRS - 2012	13.3	
Estimated Live Births	19,775	
Reported Infant Deaths	5	
Infant deaths per 1000 Estimated live Births	0	
Early Neonatal deaths per 1000 Estimated live Births	0.10	
Late Neonatal deaths per 1000 Estimated live Births	0.00	
	Total	%ge against Total Infant Deaths reported
Number of cases of Infant deaths within 24 hrs of birth	-	0%

## Goa - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	2	40%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	-	0%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	-	0%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	-	0%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	-	0%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	-	0%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	-	0%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	-	0%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	1	20%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	-	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	1	20%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	-	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	1	20%

Gujarat - Causes of Infant Deaths - HMIS - 2012 - 13		
Estimated Population - 2012	61,460,863	
CBR - SRS - 2012	21.3	
Estimated Live Births	1,309,116	
Reported Infant Deaths	15,375	
Infant deaths per 1000 Estimated live Births	12	
Early Neonatal deaths per 1000 Estimated live Births	7.57	
Late Neonatal deaths per 1000 Estimated live Births	1.77	
	Total	%ge against Total Infant Deaths reported
Number of cases of Infant deaths within 24 hrs of birth	1,734	11%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	224	1%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	164	1%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	893	6%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	224	1%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	3,505	23%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	793	5%

Gujarat - Causes of Infant Deaths - HMIS - 2012 - 13		
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	3,560	23%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	1,132	7%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	164	1%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	208	1%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	75	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	9	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	2,690	17%

Haryana - Causes of Infant Deaths - HMIS - 2012 - 13	
Estimated Population - 2012	25,834,790
CBR - SRS - 2012	21.8
Estimated Live Births	563,198
Reported Infant Deaths	4,270
Infant deaths per 1000 Estimated live Births	8

## Haryana - Causes of Infant Deaths - HMIS - 2012 - 13

Early Neonatal deaths per 1000 Estimated live Births	3.66	
Late Neonatal deaths per 1000 Estimated live Births	1.67	
	<b>Total</b>	<b>%ge against Total Infant Deaths reported</b>
Number of cases of Infant deaths within 24 hrs of birth	254	6%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	105	2%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	59	1%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	205	5%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	105	2%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	262	6%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	123	3%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	1,235	29%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	651	15%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	167	4%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	68	2%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever	63	1%



## Haryana - Causes of Infant Deaths - HMIS - 2012 - 13

related		
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	-	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	973	23%

Himachal Pradesh - Causes of Infant Deaths - HMIS - 2012 - 13		
Estimated Population - 2012	6,938,787	
CBR - SRS - 2012	16.5	
Estimated Live Births	114,490	
Reported Infant Deaths	957	
Infant deaths per 1000 Estimated live Births	8	
Early Neonatal deaths per 1000 Estimated live Births	5.00	
Late Neonatal deaths per 1000 Estimated live Births	1.31	
	Total	%ge against Total Infant Deaths reported
Number of cases of Infant deaths within 24 hrs of birth	244	25%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	51	5%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	27	3%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	79	8%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	17	2%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	78	8%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	24	3%

### Himachal Pradesh - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	121	13%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	82	9%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	70	7%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	19	2%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	38	4%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	-	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	107	11%

### Jammu & Kashmir - Causes of Infant Deaths - HMIS - 2012 - 13

Estimated Population - 2012	12,825,002
CBR - SRS - 2012	17.8
Estimated Live Births	228,285
Reported Infant Deaths	76
Infant deaths per 1000 Estimated live Births	0
Early Neonatal deaths per 1000 Estimated live Births	0.15

## Jammu & Kashmir - Causes of Infant Deaths - HMIS - 2012 - 13

Late Neonatal deaths per 1000 Estimated live Births	0.02	
	Total	%ge against Total Infant Deaths reported
Number of cases of Infant deaths within 24 hrs of birth	6	8%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	7	9%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	3	4%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	7	9%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	1	1%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	6	8%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	1	1%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	8	11%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	-	0%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	6	8%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	1	1%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	4	5%

### Jammu & Kashmir - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	-	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	26	34%

### Jharkhand - Causes of Infant Deaths - HMIS - 2012 - 13

Estimated Population - 2012	33,642,316	
CBR - SRS - 2012	25	
Estimated Live Births	841,058	
Reported Infant Deaths	1,108	
Infant deaths per 1000 Estimated live Births	1	
Early Neonatal deaths per 1000 Estimated live Births	0.71	
Late Neonatal deaths per 1000 Estimated live Births	0.11	
	<b>Total</b>	<b>%ge against Total Infant Deaths reported</b>
Number of cases of Infant deaths within 24 hrs of birth	261	24%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	24	2%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	2	0%

### Jharkhand - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	29	3%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	3	0%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	117	11%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	29	3%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	165	15%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	62	6%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	51	5%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	13	1%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	54	5%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	-	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	298	27%

Karnataka - Causes of Infant Deaths - HMIS - 2012 - 13		
Estimated Population - 2012	62,040,644	
CBR - SRS - 2012	18.8	
Estimated Live Births	1,166,364	
Reported Infant Deaths	3,237	
Infant deaths per 1000 Estimated live Births	3	
Early Neonatal deaths per 1000 Estimated live Births	1.49	
Late Neonatal deaths per 1000 Estimated live Births	0.47	
	Total	%ge against Total Infant Deaths reported
Number of cases of Infant deaths within 24 hrs of birth	500	15%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	190	6%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	96	3%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	371	11%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	93	3%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	405	13%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	191	6%

### Karnataka - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	272	8%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	174	5%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	229	7%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	26	1%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	70	2%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	3	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	617	19%

### Kerala - Causes of Infant Deaths - HMIS - 2012 - 13

Estimated Population - 2012	33,546,669
CBR - SRS - 2012	15.2
Estimated Live Births	509,909
Reported Infant Deaths	1,104
Infant deaths per 1000 Estimated live Births	2
Early Neonatal deaths per 1000 Estimated live Births	1.57



### Kerala - Causes of Infant Deaths - HMIS - 2012 - 13

Late Neonatal deaths per 1000 Estimated live Births	0.33	
	<b>Total</b>	<b>%ge against Total Infant Deaths reported</b>
Number of cases of Infant deaths within 24 hrs of birth	231	21%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	19	2%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	12	1%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	114	10%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	16	1%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	222	20%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	67	6%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	214	19%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	72	7%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	25	2%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	-	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	4	0%

### Kerala - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	-	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	108	10%

### Madhya Pradesh - Causes of Infant Deaths - HMIS - 2012 - 13

Estimated Population - 2012	73,947,121	
CBR - SRS - 2012	26.9	
Estimated Live Births	1,989,178	
Reported Infant Deaths	14,764	
Infant deaths per 1000 Estimated live Births	7	
Early Neonatal deaths per 1000 Estimated live Births	4.67	
Late Neonatal deaths per 1000 Estimated live Births	1.29	
	<b>Total</b>	<b>%ge against Total Infant Deaths reported</b>
Number of cases of Infant deaths within 24 hrs of birth	3,462	23%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	485	3%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	250	2%

## Madhya Pradesh - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	864	6%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	252	2%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	1,627	11%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	530	4%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	2,846	19%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	1,536	10%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	535	4%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	99	1%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	212	1%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	13	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	2,053	14%

Maharashtra - Causes of Infant Deaths - HMIS - 2012 - 13		
Estimated Population - 2012	114,067,506	
CBR - SRS - 2012	16.7	
Estimated Live Births	1,904,927	
Reported Infant Deaths	34,357	
Infant deaths per 1000 Estimated live Births	18	
Early Neonatal deaths per 1000 Estimated live Births	12.07	
Late Neonatal deaths per 1000 Estimated live Births	2.63	
	Total	%ge against Total Infant Deaths reported
Number of cases of Infant deaths within 24 hrs of birth	7,285	21%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	1,138	3%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	572	2%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	3,207	9%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	530	2%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	6,560	19%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	1,531	4%

### Maharashtra - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	4,809	14%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	2,386	7%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	1,198	3%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	225	1%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	391	1%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	31	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	4,494	13%

### Manipur - Causes of Infant Deaths - HMIS - 2012 - 13

Estimated Population - 2012	2,757,523
CBR - SRS - 2012	14.4
Estimated Live Births	39,708
Reported Infant Deaths	43
Infant deaths per 1000 Estimated live Births	1
Early Neonatal deaths per 1000 Estimated live Births	0.40

## Manipur - Causes of Infant Deaths - HMIS - 2012 - 13

Late Neonatal deaths per 1000 Estimated live Births	0.08	
	Total	%ge against Total Infant Deaths reported
Number of cases of Infant deaths within 24 hrs of birth	5	12%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	2	5%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	-	0%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	2	5%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	-	0%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	4	9%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	-	0%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	3	7%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	3	7%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	7	16%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	1	2%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	6	14%

### Manipur - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	-	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	10	23%

### Meghalaya - Causes of Infant Deaths - HMIS - 2012 - 13

Estimated Population - 2012	3,039,344	
CBR - SRS - 2012	24.1	
Estimated Live Births	73,248	
Reported Infant Deaths	2,200	
Infant deaths per 1000 Estimated live Births	30	
Early Neonatal deaths per 1000 Estimated live Births	13.47	
Late Neonatal deaths per 1000 Estimated live Births	1.27	
	<b>Total</b>	<b>%ge against Total Infant Deaths reported</b>
Number of cases of Infant deaths within 24 hrs of birth	441	20%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	82	4%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	3	0%

### Meghalaya - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	108	5%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	3	0%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	95	4%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	30	1%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	261	12%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	57	3%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	246	11%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	69	3%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	219	10%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	1	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	585	27%



Mizoram - Causes of Infant Deaths - HMIS - 2012 - 13		
Estimated Population - 2012	1,113,327	
CBR - SRS - 2012	16.6	
Estimated Live Births	18,481	
Reported Infant Deaths	621	
Infant deaths per 1000 Estimated live Births	34	
Early Neonatal deaths per 1000 Estimated live Births	6.38	
Late Neonatal deaths per 1000 Estimated live Births	0.54	
	Total	%ge against Total Infant Deaths reported
Number of cases of Infant deaths within 24 hrs of birth	9	1%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	32	5%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	5	1%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	14	2%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	1	0%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	33	5%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	2	0%

### Mizoram - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	30	5%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	2	0%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	215	35%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	11	2%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	78	13%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	-	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	189	30%

### Nagaland - Causes of Infant Deaths - HMIS - 2012 - 13

Estimated Population - 2012	1,990,505
CBR - SRS - 2012	16.1
Estimated Live Births	32,047
Reported Infant Deaths	74
Infant deaths per 1000 Estimated live Births	2

Nagaland - Causes of Infant Deaths - HMIS - 2012 - 13		
Early Neonatal deaths per 1000 Estimated live Births	0.94	
Late Neonatal deaths per 1000 Estimated live Births	0.25	
	Total	%ge against Total Infant Deaths reported
Number of cases of Infant deaths within 24 hrs of birth	4	5%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	4	5%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	2	3%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	9	12%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	2	3%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	4	5%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	-	0%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	9	12%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	4	5%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	12	16%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	6	8%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever	7	9%

### Nagaland - Causes of Infant Deaths - HMIS - 2012 - 13

related		
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	-	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	11	15%

### Odisha - Causes of Infant Deaths - HMIS - 2012 - 13

Estimated Population - 2012	42,510,898	
CBR - SRS - 2012	20.1	
Estimated Live Births	854,469	
Reported Infant Deaths	11,102	
Infant deaths per 1000 Estimated live Births	13	
Early Neonatal deaths per 1000 Estimated live Births	7.17	
Late Neonatal deaths per 1000 Estimated live Births	2.59	
	Total	%ge against Total Infant Deaths reported
Number of cases of Infant deaths within 24 hrs of birth	1,255	11%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	252	2%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	198	2%

### Odisha - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	1,154	10%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	186	2%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	2,084	19%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	983	9%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	1,383	12%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	844	8%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	886	8%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	46	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	90	1%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	6	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	1,735	16%

Punjab - Causes of Infant Deaths - HMIS - 2012 - 13		
Estimated Population - 2012	28,064,391	
CBR - SRS - 2012	16.2	
Estimated Live Births	454,643	
Reported Infant Deaths	2,106	
Infant deaths per 1000 Estimated live Births	5	
Early Neonatal deaths per 1000 Estimated live Births	2.58	
Late Neonatal deaths per 1000 Estimated live Births	0.77	
	Total	%ge against Total Infant Deaths reported
Number of cases of Infant deaths within 24 hrs of birth	469	22%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	48	2%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	18	1%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	82	4%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	12	1%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	98	5%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	37	2%

### Punjab - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	474	23%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	281	13%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	99	5%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	30	1%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	78	4%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	-	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	380	18%

## Rajasthan - Causes of Infant Deaths - HMIS - 2012 - 13

Estimated Population - 2012	69,971,092	
CBR - SRS - 2012	26.2	
Estimated Live Births	1,833,243	
Reported Infant Deaths	10,961	
Infant deaths per 1000 Estimated live Births	6	
Early Neonatal deaths per 1000 Estimated live Births	3.85	
Late Neonatal deaths per 1000 Estimated live Births	0.85	
	Total	%ge against Total Infant Deaths reported
Number of cases of Infant deaths within 24 hrs of birth	1,806	16%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	447	4%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	194	2%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	1,050	10%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	179	2%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	1,559	14%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	395	4%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and	2,203	20%



Rajasthan - Causes of Infant Deaths - HMIS - 2012 - 13		
LBW		
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	795	7%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	229	2%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	265	2%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	485	4%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	1	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	1,353	12%

Tripura - Causes of Infant Deaths - HMIS - 2012 - 13	
Estimated Population - 2012	3,722,820
CBR - SRS - 2012	14.3
Estimated Live Births	53,236
Reported Infant Deaths	280
Infant deaths per 1000 Estimated live Births	5
Early Neonatal deaths per 1000 Estimated live Births	1.80

### Tripura - Causes of Infant Deaths - HMIS - 2012 - 13

Late Neonatal deaths per 1000 Estimated live Births	0.68	
	<b>Total</b>	<b>%ge against Total Infant Deaths reported</b>
Number of cases of Infant deaths within 24 hrs of birth	13	5%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	20	7%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	19	7%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	24	9%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	5	2%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	20	7%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	5	2%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	19	7%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	7	3%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	61	22%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	2	1%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	8	3%

### Tripura - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	1	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	76	27%

### Uttarakhand - Causes of Infant Deaths - HMIS - 2012 - 13

Estimated Population - 2012	10,296,933	
CBR - SRS - 2012	18.9	
Estimated Live Births	194,612	
Reported Infant Deaths	72	
Infant deaths per 1000 Estimated live Births	0	
Early Neonatal deaths per 1000 Estimated live Births	0.20	
Late Neonatal deaths per 1000 Estimated live Births	0.03	
	Total	%ge against Total Infant Deaths reported
Number of cases of Infant deaths within 24 hrs of birth	20	28%

### Uttarakhand - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	3	4%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	4	6%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	1	1%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	-	0%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	7	10%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	-	0%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	8	11%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	2	3%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	9	13%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	4	6%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	5	7%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	-	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	9	13%

Uttar Pradesh - Causes of Infant Deaths - HMIS - 2012 - 13		
Estimated Population - 2012	203,286,070	
CBR - SRS - 2012	27.8	
Estimated Live Births	5,651,353	
Reported Infant Deaths	1,011	
Infant deaths per 1000 Estimated live Births	0	
Early Neonatal deaths per 1000 Estimated live Births	0.09	
Late Neonatal deaths per 1000 Estimated live Births	0.04	
	Total	%ge against Total Infant Deaths reported
Number of cases of Infant deaths within 24 hrs of birth	29	3%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	58	6%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	30	3%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	86	9%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	22	2%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	142	14%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	40	4%

### Uttar Pradesh - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	189	19%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	115	11%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	51	5%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	37	4%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	81	8%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	11	1%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	120	12%

### West Bengal - Causes of Infant Deaths - HMIS - 2012 - 13

Estimated Population - 2012	92,541,954
CBR - SRS - 2012	16.3
Estimated Live Births	1,508,434
Reported Infant Deaths	15,478
Infant deaths per 1000 Estimated live Births	10
Early Neonatal deaths per 1000 Estimated live Births	6.87

## West Bengal - Causes of Infant Deaths - HMIS - 2012 - 13

Late Neonatal deaths per 1000 Estimated live Births	1.32	
	<b>Total</b>	<b>%ge against Total Infant Deaths reported</b>
Number of cases of Infant deaths within 24 hrs of birth	1,450	9%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Sepsis	1,160	7%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Sepsis	483	3%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being Asphyxia	2,761	18%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being Asphyxia	211	1%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being LBW	2,347	15%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being LBW	788	5%
Number of cases of Infant deaths 24 hours to one week of birth with the probable cause being other than Sepsis, Asphyxia and LBW	2,644	17%
Number of cases of Infant deaths one week to 4 weeks of birth with the probable cause being other than Sepsis, Asphyxia and LBW	516	3%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Pneumonia	723	5%
Number of cases of Infant or Child deaths between 1 month to 11 months of age with the probable cause being Diarrhoea	46	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Fever related	292	2%

### West Bengal - Causes of Infant Deaths - HMIS - 2012 - 13

Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being Measles	3	0%
Number of cases of Infant or Child deaths between 1 month to 11 month of age with the probable cause being other than Pneumonia, Diarrhoea, Fever related and Measles	2,054	13%